
 Scheda di frequenza 1 sessione 2, metodo: Mitchell, 09 Ott 2005
 Neuberg = S

	1	2	3	4	5	6	7	8	9	10	11	12
A :	130	450	920	-50	450	500	420	420	300	450	450	170
B :	450	420	170	450	450	920	-150	450	150	-150	420	420
C :	450	450	-50	420	920	-150	16060	450	170	450	-50	170

score	freq	NS	EW
920	3	67.91	2.09
500	1	63.80	6.20
450	12	50.43	19.57
420	6	31.91	38.09
300	1	24.71	45.29
170	4	19.57	50.43
150	1	14.43	55.57
130	1	12.37	57.63
-50	3	8.26	61.74
-150	3	2.09	67.91

scores 35 + 1 attrib
 totsco 36

top = 68, integ = 70
 N.fac = 1.02857, N.add = 0.0286

 Scheda di frequenza 10 sessione 2, metodo: Mitchell, 09 Ott 2005
 Neuberg = S

	1	2	3	4	5	6	7	8	9	10	11	12
A :	200	100	100	200	200	-120	100	100	-90	-800	-90	300
B :	200	100	-120	300	-90	100	100	200	100	100	100	200
C :	100	100	100	200	400	-150	200	100	16060	200	100	-90

score	freq	NS	EW
400	1	69.97	0.03
300	2	66.89	3.11
200	9	55.57	14.43
100	15	30.89	39.11
-90	4	11.34	58.66
-120	2	5.17	64.83
-150	1	2.09	67.91
-800	1	0.03	69.97

scores 35 + 1 attrib
 totsco 36

top = 68, integ = 70
 N.fac = 1.02857, N.add = 0.0286

 Scheda di frequenza 11 sessione 2, metodo: Mitchell, 09 Ott 2005
 Neuberg = S

	1	2	3	4	5	6	7	8	9	10	11	12
A :	-420	50	-170	-200	14060	-170	-170	-300	-420	-420	-420	-420
B :	-590	-420	-170	-170	-500	-420	-420	-420	-420	-420	-420	-170
C :	-420	-420	16060	-140	-420	-420	-140	-420	-800	-420	-420	-200

score	freq	NS	EW
50	1	69.94	0.06
-140	2	66.76	3.24
-170	6	58.29	11.71
-200	2	49.82	20.18
-300	1	46.65	23.35
-420	19	25.47	44.53
-500	1	4.29	65.71
-590	1	2.18	67.82
-800	1	0.06	69.94

scores 34 + 2 attrib
 totsco 36

top = 66, integ = 70
 N.fac = 1.05882, N.add = 0.0588

 Scheda di frequenza 12 sessione 2, metodo: Mitchell, 09 Ott 2005
 Neuberg = S

	1	2	3	4	5	6	7	8	9	10	11	12
A :	-420	-420	-420	50	-450	-480	50	-100	-420	-200	50	50
B :	-420	100	-420	-420	-140	-590	50	-170	-480	50	50	-480
C :	-420	50	16060	50	-420	50	50	-420	-420	50	-420	100

score	freq	NS	EW
100	2	68.94	1.06
50	12	54.54	15.46
-100	1	41.17	28.83
-140	1	39.11	30.89
-170	1	37.06	32.94
-200	1	35.00	35.00
-420	12	21.63	48.37
-450	1	8.26	61.74
-480	3	4.14	65.86
-590	1	0.03	69.97

scores 35 + 1 attrib
 totsco 36

top = 68, integ = 70
 N.fac = 1.02857, N.add = 0.0286

 Scheda di frequenza 13 sessione 2, metodo: Mitchell, 09 Ott 2005
 Neuberg = S

	1	2	3	4	5	6	7	8	9	10	11	12
A :	110	600	-100	870	600	600	-100	600	-100	-100	630	-100
B :	-100	720	-100	-200	-300	190	-100	660	-100	620	-100	620
C :	-200	690	-100	600	800	630	150	690	720	16060	600	-100

score	freq	NS	EW
870	1	69.97	0.03
800	1	67.91	2.09
720	2	64.83	5.17
690	2	60.71	9.29
660	1	57.63	12.37
630	2	54.54	15.46
620	2	50.43	19.57
600	6	42.20	27.80
190	1	35.00	35.00
150	1	32.94	37.06
110	1	30.89	39.11
-100	12	17.51	52.49
-200	2	3.11	66.89
-300	1	0.03	69.97

scores 35 + 1 attrib
 totsco 36

top = 68, integ = 70
 N.fac = 1.02857, N.add = 0.0286

 Scheda di frequenza 14 sessione 2, metodo: Mitchell, 09 Ott 2005
 Neuberg = S

	1	2	3	4	5	6	7	8	9	10	11	12
A :	-130	500	-130	-50	-400	-570	-130	-130	-50	-130	-500	110
B :	-300	110	100	100	-130	-110	-300	-200	-50	-150	-130	-110
C :	-100	-130	-100	100	14060	50	-130	-510	-300	16060	-130	-670

score	freq	NS	EW
500	1	69.94	0.06
110	2	66.76	3.24
100	3	61.47	8.53
50	1	57.24	12.76
-50	3	53.00	17.00
-100	2	47.71	22.29
-110	2	43.47	26.53
-130	10	30.76	39.24
-150	1	19.12	50.88
-200	1	17.00	53.00
-300	3	12.76	57.24
-400	1	8.53	61.47
-500	1	6.41	63.59
-510	1	4.29	65.71
-570	1	2.18	67.82
-670	1	0.06	69.94

scores 34 + 2 attrib
 totsco 36

top = 66, integ = 70
 N.fac = 1.05882, N.add = 0.0588

 Scheda di frequenza 15 sessione 2, metodo: Mitchell, 09 Ott 2005
 Neuberg = S

	1	2	3	4	5	6	7	8	9	10	11	12
A :	170	50	100	-90	50	150	100	-400	110	650	-400	50
B :	-400	50	300	170	100	-90	250	110	14060	-110	100	110
C :	50	140	100	16060	50	170	100	800	170	100	100	-180

score	freq	NS	EW
800	1	69.94	0.06
650	1	67.82	2.18
300	1	65.71	4.29
250	1	63.59	6.41
170	4	58.29	11.71
150	1	53.00	17.00
140	1	50.88	19.12
110	3	46.65	23.35
100	8	35.00	35.00
50	6	20.18	49.82
-90	2	11.71	58.29
-110	1	8.53	61.47
-180	1	6.41	63.59
-400	3	2.18	67.82

scores 34 + 2 attrib
 totsco 36

top = 66, integ = 70
 N.fac = 1.05882, N.add = 0.0588

Scheda di frequenza 16 sessione 2, metodo: Mitchell, 09 Ott 2005
Neuberg = S

1 2 3 4 5 6 7 8 9 10 11 12

A : -1430 -1430 -680 -710 -1430 -710 -1430 -1430 -620 -710 -1430 -1460
B : -1460 -1430 -680 -680 -680 -680 -710 -680 -680 -650 -640 -680
C : -1430 -680 -680 16060 -1430 -680 -680 -1430 -680 -1430 -1460 -680

score	freq	NS	EW
-620	1	69.97	0.03
-640	1	67.91	2.09
-650	1	65.86	4.14
-680	14	50.43	19.57
-710	4	31.91	38.09
-1430	11	16.49	53.51
-1460	3	2.09	67.91

scores 35 + 1 attrib
totsco 36

top = 68, integ = 70
N.fac = 1.02857, N.add = 0.0286

 Scheda di frequenza 17 sessione 2, metodo: Mitchell, 09 Ott 2005
 Neuberg = S

	1	2	3	4	5	6	7	8	9	10	11	12
A :	50	50	100	-400	50	-300	550	100	100	-400	-140	140
B :	50	-100	-50	110	800	150	590	100	100	100	-300	170
C :	-100	-50	-500	-140	-100	300	50	-170	-210	100	16060	50

score	freq	NS	EW
800	1	69.97	0.03
590	1	67.91	2.09
550	1	65.86	4.14
300	1	63.80	6.20
170	1	61.74	8.26
150	1	59.69	10.31
140	1	57.63	12.37
110	1	55.57	14.43
100	7	47.34	22.66
50	6	33.97	36.03
-50	2	25.74	44.26
-100	3	20.60	49.40
-140	2	15.46	54.54
-170	1	12.37	57.63
-210	1	10.31	59.69
-300	2	7.23	62.77
-400	2	3.11	66.89
-500	1	0.03	69.97

scores 35 + 1 attrib
 totsco 36

top = 68, integ = 70
 N.fac = 1.02857, N.add = 0.0286

 Scheda di frequenza 18 sessione 2, metodo: Mitchell, 09 Ott 2005
 Neuberg = S

	1	2	3	4	5	6	7	8	9	10	11	12
A :	630	660	210	-200	600	660	630	630	660	630	600	630
B :	600	660	660	630	150	630	600	660	600	630	600	660
C :	630	660	630	630	660	630	600	600	630	-100	16060	600

score	freq	NS	EW
660	9	61.74	8.26
630	13	39.11	30.89
600	9	16.49	53.51
210	1	6.20	63.80
150	1	4.14	65.86
-100	1	2.09	67.91
-200	1	0.03	69.97

scores 35 + 1 attrib
 totsco 36

top = 68, integ = 70
 N.fac = 1.02857, N.add = 0.0286

 Scheda di frequenza 19 sessione 2, metodo: Mitchell, 09 Ott 2005
 Neuberg = S

	1	2	3	4	5	6	7	8	9	10	11	12
A :	-50	-50	-110	110	-100	-50	-100	110	-50	130	-50	110
B :	130	-50	150	-250	110	-50	120	-300	130	130	-50	-110
C :	-50	120	90	-90	16060	-50	460	200	110	430	90	-50

score	freq	NS	EW
460	1	69.97	0.03
430	1	67.91	2.09
200	1	65.86	4.14
150	1	63.80	6.20
130	4	58.66	11.34
120	2	52.49	17.51
110	5	45.29	24.71
90	2	38.09	31.91
-50	11	24.71	45.29
-90	1	12.37	57.63
-100	2	9.29	60.71
-110	2	5.17	64.83
-250	1	2.09	67.91
-300	1	0.03	69.97

scores 35 + 1 attrib
 totsco 36

top = 68, integ = 70
 N.fac = 1.02857, N.add = 0.0286

 Scheda di frequenza 21 sessione 2, metodo: Mitchell, 09 Ott 2005
 Neuberg = S

	1	2	3	4	5	6	7	8	9	10	11	12
A :	-400	-400	-400	50	-400	-400	-400	50	-400	50	-90	50
B :	50	-110	-130	-150	50	50	-400	-400	50	-430	-100	50
C :	-400	-120	-400	-400	-430	100	-400	-430	-400	50	-400	16060

score	freq	NS	EW
100	1	69.97	0.03
50	10	58.66	11.34
-90	1	47.34	22.66
-100	1	45.29	24.71
-110	1	43.23	26.77
-120	1	41.17	28.83
-130	1	39.11	30.89
-150	1	37.06	32.94
-400	15	20.60	49.40
-430	3	2.09	67.91

scores 35 + 1 attrib
 totsco 36

top = 68, integ = 70
 N.fac = 1.02857, N.add = 0.0286

 Scheda di frequenza 22 sessione 2, metodo: Mitchell, 09 Ott 2005
 Neuberg = S

	1	2	3	4	5	6	7	8	9	10	11	12
A :	120	-100	100	150	180	110	460	400	800	130	210	110
B :	130	120	150	130	110	150	430	430	430	130	150	110
C :	150	210	400	430	110	400	120	150	110	400	-50	16060

score	freq	NS	EW
800	1	69.97	0.03
460	1	67.91	2.09
430	4	62.77	7.23
400	4	54.54	15.46
210	2	48.37	21.63
180	1	45.29	24.71
150	6	38.09	31.91
130	4	27.80	42.20
120	3	20.60	49.40
110	6	11.34	58.66
100	1	4.14	65.86
-50	1	2.09	67.91
-100	1	0.03	69.97

scores 35 + 1 attrib
 totsco 36

top = 68, integ = 70
 N.fac = 1.02857, N.add = 0.0286

 Scheda di frequenza 23 sessione 2, metodo: Mitchell, 09 Ott 2005
 Neuberg = S

	1	2	3	4	5	6	7	8	9	10	11	12
A :	100	100	100	100	-600	300	200	-630	670	-630	100	100
B :	-630	670	200	-600	100	-600	300	-800	-600	360	100	100
C :	100	-630	100	100	-500	16060	-100	-130	100	-660	-600	100

score	freq	NS	EW
670	2	68.94	1.06
360	1	65.86	4.14
300	2	62.77	7.23
200	2	58.66	11.34
100	14	42.20	27.80
-100	1	26.77	43.23
-130	1	24.71	45.29
-500	1	22.66	47.34
-600	5	16.49	53.51
-630	4	7.23	62.77
-660	1	2.09	67.91
-800	1	0.03	69.97

scores 35 + 1 attrib
 totsco 36

top = 68, integ = 70
 N.fac = 1.02857, N.add = 0.0286

 Scheda di frequenza 24 sessione 2, metodo: Mitchell, 09 Ott 2005
 Neuberg = S

	1	2	3	4	5	6	7	8	9	10	11	12
A :	170	-50	500	170	50	300	-50	100	650	800	170	420
B :	420	590	590	170	420	170	170	170	450	-50	690	140
C :	300	-610	590	-100	300	16060	100	170	450	420	100	-110

score	freq	NS	EW
800	1	69.97	0.03
690	1	67.91	2.09
650	1	65.86	4.14
590	3	61.74	8.26
500	1	57.63	12.37
450	2	54.54	15.46
420	4	48.37	21.63
300	3	41.17	28.83
170	8	29.86	40.14
140	1	20.60	49.40
100	3	16.49	53.51
50	1	12.37	57.63
-50	3	8.26	61.74
-100	1	4.14	65.86
-110	1	2.09	67.91
-610	1	0.03	69.97

scores 35 + 1 attrib
 totsco 36

top = 68, integ = 70
 N.fac = 1.02857, N.add = 0.0286

 Scheda di frequenza 25 sessione 2, metodo: Mitchell, 09 Ott 2005
 Neuberg = S

	1	2	3	4	5	6	7	8	9	10	11	12
A :	-620	-650	-650	-620	-620	-620	-500	-650	-620	-620	-620	-170
B :	-650	-620	-620	110	-650	-170	-650	-620	-620	-100	-620	-620
C :	100	-620	-620	-620	-620	-650	200	-620	-650	100	-620	-170

score	freq	NS	EW
200	1	70.00	0.00
110	1	68.00	2.00
100	2	65.00	5.00
-100	1	62.00	8.00
-170	3	58.00	12.00
-500	1	54.00	16.00
-620	19	34.00	36.00
-650	8	7.00	63.00

scores 36
 totsco 36

top = 70
 N.fac = 1.00000, N.add = 0.0000

